

**REGISTRATION FOR SPECIAL LOCAL NEED  
FOR DISTRIBUTION AND USE ONLY WITHIN MONTANA**



**EMULSIFIABLE CONCENTRATE FOR AGRICULTURAL USE ONLY**  
EPA Reg. No. 10163-277

<b>ACTIVE INGREDIENT:</b> Hexythiazox	<b>% By Wt.</b>
<i>trans</i> -5-(4-Chlorophenyl)-N-cyclohexyl-4-methyl-2-oxothiazolidine-3-carboxamide .....	11.8%
<b>OTHER INGREDIENTS</b> .....	<u>88.2%</u>
	<b>TOTAL 100.00%</b>

Onager contains 1.0 lb. active ingredient per gallon  
Contains Petroleum Distillate

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- All applicable directions, restrictions and precautions on the USEPA registered label are to be followed.
- This labeling must be in the possession of the user at the time of pesticide application.

CROP	PEST	RATE OzS./Acre	COMMENTS
<b>ALFALFA GROWN FOR SEED</b>	Twospotted Spider Mite, Pacific Spider Mite, Strawberry Spider Mite	12 – 16	Onager is active against all immature stages of Tetranychid spider mites, including freshly laid eggs. Therefore, it is important to apply Onager prior to adult mite buildup. Scout and apply early when mites first begin colonizing the upper $\frac{1}{3}$ to $\frac{1}{2}$ of the plant canopy. An early application timing and thorough coverage are essential for acceptable control. Apply Onager by ground in a minimum of 20 gallons of water per acre at a minimum of 35 psi to insure optimum coverage.
	<ul style="list-style-type: none"> <li>• Do not cut current year's treated seed crop for hay or forage.</li> <li>• Do not graze current year's treated seed crop.</li> <li>• Screenings from seed processing are prohibited from food or feed channels. All Onager treated seed screenings must be removed from the food or feed market.</li> <li>• Treated seed is not to be used for sprouting. All seed treated with Onager is to be tagged at processing plants "<b>NOT FOR HUMAN OR ANIMAL CONSUMPTION</b>". It shall be the grower's responsibility to notify the processing plants of any seed crop treated with Onager.</li> <li>• Keep a minimum distance of 200 feet from the edge of an alfalfa field to the center of a stream</li> </ul>		

**RESTRICTIONS**

- Do not enter treated area without label prescribed Personal Protective Equipment for 12 hours after application.
- Livestock shall not be allowed to graze or feed on current years treated alfalfa seed crop.
- The crop refuse and seed cleanings from current years treated alfalfa seed crop cannot be used for pasture, feed, food, forage, or bedding purposes.
- Alfalfa seed from treated fields may not be used for growing sprouts for human or animal consumption.
- Onager is non-hazardous to bees and may be applied during bloom. However, if tank mixed with another product that may be hazardous to bees, the tank mix spray should not be applied when bees are actively visiting the treatment area. Follow label restrictions of any tank mix product.
- Refer to the federally registered label for precautionary statements and statement of practical treatment.
- Do not apply within 4 weeks of harvest.
- Do not make more than one application per growing season.

**24(c) REGISTRANT**

Gowan Company  
PO Box 5569  
Yuma, AZ 85366