



Dow AgroSciences

Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

FIFRA 24(c) Special Local Need Label (SLN)

For Distribution and Use Only in the State of Montana
Control of Spotted Wing Drosophila in Cherry

Entrust[®] SC

EPA Reg. No. 62719-621

SLN MT -12-0004

This label expires and must not be distributed or used in accordance with this SLN registration after

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This state-specific Section 24(c) labeling must be in the possession of the user at the time of application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

Refer to product label for Entrust SC for Use Precautions, Mixing Directions, and Application Directions.

Application Timing: Treat when pests appears and crop is becoming susceptible (e.g., around the blush or pink stage) to spotted wing drosophila (SWD). Fruits become more susceptible as harvest approaches. Later harvested cultivars are at greater risk from SWD.

Application Rate: Apply 4.8 to 6.4 fl oz of Entrust SC per acre as a foliar spray to control spotted wing drosophila. Use a higher rate in the rate range for moderate to severe pest infestations. Heavy infestations may require repeat applications, but follow resistance management guidelines.

Resistance Management: Do not make more than 2 consecutive applications of Group 5 insecticides (spinosad and spinetoram). If additional applications are required, rotate to an effective product in a different group for at least 2 applications. Do not make more than three applications per acre per year of Group 5 insecticides (spinetoram [Delegate[®] WG] and spinosad [Entrust[®], Entrust[®] SC Success[®]]).

Treatment Schedule: If three Entrust SC applications are made, the first may be made up to 28 days before harvest, the second up to 10 days before harvest, and the final application at 3 days before harvest. Two applications of one or more effective insecticides not in Group 5 must be made between the first and second Entrust SC applications.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 3 days of harvest.
- Allow at least 7 days between applications.
- Do not apply more than 6.4 fl oz of Entrust SC (0.10 lb ai) per acre per application.
- Do not apply more than a total of 12.8 fl oz of Entrust SC (0.20 lb ai) per acre within 10 days of harvest.
- Do not apply more than a total of 19.2 fl oz of Entrust SC (0.30 lb ai) per acre per year.
- Apply by air blast sprayer. Do not use hand held or backpack sprayers.
- **Chemigation:** For use under this SLN label, do not apply this product through any type of irrigation system.

Dow AgroSciences LLC
9330 Zionsville Road
Indianapolis, Indiana 46268
1- 800-992-5994 (option #4)