

DIRECTIONS FOR USE

For use only against Columbian and Richardson ground squirrels.

Do not apply on roads, near residential or public use areas or over water.

Apply bait only once per season at a time when ground squirrels show good acceptance of grains.

Do not apply if rain is forecast within 2 days after bait application.

HAND BAITING

Apply bait only to bare ground within 3 ft of ground squirrel burrow entrances in rangeland, pastures, alfalfa, wheat, oats, barley and adjacent noncrop areas.

Pretreatment: Do not apply bait unless tests with untreated oats show good acceptance of grains by ground squirrels in areas to be treated.

Prebaiting (Strongly Recommended): Scatter one teaspoon of nontoxic oats on bare ground near all active ground squirrel burrows in the treatment area. Do not apply toxic bait until nontoxic prebait is consumed. Prebaiting has been shown to increase acceptance of toxic bait by the target ground squirrel species.

Baiting: Use an appropriately calibrated bait dipper to apply bait at 4 grams (one teaspoonful) per active burrow.

Scatter bait evenly on bare ground near burrow entrance. Do not place bait in piles or pour down the burrow entrance.

RESTRICTED USE PESTICIDE

For retail sale and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

ZP[®]

Rodent Oat Bait AG

For distribution and use in MONTANA only. This label must be in the possession of the user at the time of application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. All applicable directions, restrictions and precautions on the EPA registered label are to be followed.

NET CONTENTS: 45 lbs

MONTANA REGISTRATION
SEC. 24 (c) - SLN NO. MT-890009

This label is valid through December 31, 2021 or until otherwise amended, withdrawn, cancelled or suspended.

Manufactured by:



Bell Laboratories, Inc.
Madison, WI 53704 U.S.A.

EPA REG. NO. 12455-102

EPA EST. NO. 12455-WI-1

BROADCAST BAITING

Apply only to rangeland and adjacent noncrop areas. Do not broadcast bait on sites where ground cover is greater than 50 percent where squirrels will not find sufficient bait to be effective.

Pretreatment: Do not apply bait unless tests with untreated oats show good acceptance of grains by ground squirrels in the areas to be treated.

Prebaiting (Strongly Recommended): Broadcast 3 to 6 pounds of nontoxic oats per swath acre 2 to 3 days prior to applying toxic bait. Prebaiting has been shown to increase acceptance of the toxic bait by the target ground squirrel species.

Baiting: Broadcast 3 to 6 pounds of toxic bait per swath acre, not to exceed 6 pounds of bait per acre. Use the higher rates on sites with denser vegetative growth. An untreated interval varying from 20 feet but not to exceed 100 feet may be left between baited swaths depending on the density of ground cover. Apply the toxic bait swaths to the same sites that prebait was applied.