

# **Special Local Need**

### FOR USE ONLY IN THE STATE OF MONTANA

### RESTRICTED USE PESTICIDE

Restricted Use Pesticide due to carcinogenicity, potential for affecting fetal development, and high acute toxicity to humans. For retail sale to and use by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicators Certification.

## Super Tin® 80WP Fungicide

EPA Registration No. 70506-214

EPA SLN No. MT-090001 Good through 2012

### **Application Instructions for Control of Cercospora on Sugar Beet**

### **IMPORTANT:**

BEFORE USING SUPER TIN<sup>®</sup> 80WP Fungicide, READ AND CAREFULLY NOTE THE PRECAUTIONARY STATEMENTS AND OTHER PROCEDURAL INFORMATION APPEARING ON THE EPA REGISTERED LABEL OR ON OTHER SUPPLEMENTAL LABELS.

**Use Rate:** 2.5-5.0 oz/acre (15-7.5 acres/2.34 lb pack or 10-5 acres/25.0 oz. pack or 7.5-3.75 acres/18.75 oz. pack)

**Ground (Closed cabs only):** Apply in at least 15 gallons of water. Full coverage of the foliage is necessary for best results.

Aerial: (helicopter or fixed wing aircraft): Apply in 5 to 10 gallons of water.

**Ground and Aerial Applications:** Diluted spray should be directed uniformly to all parts of the plant. Use lower gallonage when plants are small and increase volume with plant size. Use the lower rate for protective sprays and the higher rates later in the season or during high infection periods. Application should begin when leaf spot conditions appear or when the disease is in the area and repeated at 10 to 14-day intervals. Do not exceed 10 ounces per acre per season. Do not treat within 7 days of harvest. Do not graze or feed beet tops to livestock.

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This label and the federally registered label must be in possession of the user at the time of pesticide application.