



Section 24(c) Special Local Need Label

BICYCLOPYRONE	GROUP	27	HERBICIDE
BROMOXYNIL	GROUP	6	HERBICIDE

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA

Talinor®

Fallow Application to Control Kochia, Russian Thistle, Palmer Amaranth and Other Difficult to Control Weeds

EPA Reg. No. 100-1570
EPA SLN No. MT-24-0002

This label expires and must not be distributed or used in accordance with this SLN registration after December 31, 2028

Active Ingredient:

Bicyclopyrone*:	3.41%
Bromoxynil Octanoate**:	23.16%
Other Ingredients:	73.43%
Total:	100.0%

*CAS No. 352010-68-5

**CAS No.1689-99-2

Talinor Herbicide is an EC formulation containing 0.31 pounds of bicyclopyrone and 1.46 pounds of bromoxynil acid equivalent per gallon.

Contains petroleum distillates.

DIRECTIONS FOR USE

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This SLN label and main EPA-registered Talinor label must be in the possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on this SLN label and the main EPA-registered Talinor label.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labelling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, or Viton ≥14 mils

For the control of broad leaf weeds, apply Talinor at a rate of 13.7-18.2 fl oz/A to the stubble of harvested crops, fallow ground, or prior to planting fall or spring cereals. Apply Talinor as a broadcast spray with either ground or aerial application equipment. Use a minimum of 10 gallons per acre by ground or 5 gallons per acre by aerial application.

Talinor may be tank-mixed with other herbicides labeled for use in fallow applications. Tank-mixing with labeled burndown herbicides such as 2,4-D, dicamba, Gramoxone SL 3.0®, or glyphosate is required for control of larger, drought-stressed, or hardened off weeds. The addition of a non-selective herbicide is required for grass weed control.

The use of CoAct+™ is not required when used as a non-crop foliar spray.

Restrictions: 1) Do not apply more than 18.2 fl oz/A of Talinor per year (0.045 lb ai/A per year) of bicyclopyrone-containing products. 2) Do not make more than one application per year. 3) Restricted Entry Interval (REI) is 24 hours.

Follow all other label directions, use restrictions and precautions, rotational guidelines, tank mixing directions on the Talinor Section 3 label.

©2024 Syngenta

Talinor®, Gramoxone®, CoAct+™ and the Syngenta logo are trademarks of a Syngenta Group Company.

24(c) Registrant:
Syngenta Crop Protection, LLC
P. O. Box 18300
Greensboro, NC 27419-8300

Label Code: MT1570054AA0424