

# FIFRA Section 24(c) Special Local Need (SLN) Label



Corteva Agriscience LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

## Stinger® HL

EPA Reg. No. 62719-747  
EPA SLN No. MT-24-0004

### Control of Broadleaf Weeds in Flax (including low linolenic varieties)

### FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF MONTANA)

This label for Stinger HL expires and must not be distributed or used in accordance with this SLN registration after December 31, 2029

#### ATTENTION

- The label and the federal label for this product must be in possession of the user at the time of pesticide application.
- Follow all applicable directions, restrictions, and precautions on the SLN label and the main EPA registered label. It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### Active Ingredient:

#### By Weight

clopyralid dimethylamine salt:

3,6-dichloro-2-pyridinecarboxylic acid dimethylammonium salt.....60.22%

Other Ingredients .....39.78%

Total .....100.00%

Acid Equivalent: clopyralid: 3,6-dichloro-2-pyridinecarboxylic acid – 48.8% (5 lb/gal)

**Keep Out of Reach of Children**

## **CAUTION**

### **Directions for Use**

Refer to product label for a complete list of weeds controlled, General Use Precautions, Mixing and Application instructions.

### **Application Timing**

Apply to flax, including low linolenic acid varieties 2 to 6 inches high and when the weeds are actively growing at rates shown in the following table. Consult the table below for additional information. Apply this product uniformly with ground or aerial equipment in 10 to 20 gallons total spray volume per acre (minimum of 5 gallons per acre by air).

<b>Target Broadleaf Weeds</b>	<b>Stinger HL (pint/acre)</b>
Canada thistle	0.2 for top growth suppression
annual sowthistle biennial wormwood dandelion dock, curly false chamomile green smartweed mayweed chamomile nightshade species sunflower wild buckwheat	0.15 - 0.2

### **Tank Mixes**

To control additional broadleaf weeds and provide effective, season long, control of perennials such as Canada thistle, this product may be applied in combination with labeled rates of a product containing MCPA ester or amine. For best results, tank mix 0.10 to 0.15 pints of Stinger HL per acre with a product containing MCPA ester or amine at a rate of 6–8 fl oz ae/A. Use the higher rates of Stinger HL and MCPA when the condition of the weeds and/or crop at the time of treatment may prevent optimum control. Apply when the majority of basal leaves have emerged from the soil (plant 4–6 inches in height) up to the bud stage. Cultivation can disrupt translocation to the roots of perennials, such as Canada thistle. For best results, do not cultivate thistle patches.

Stinger HL, alone or in tank-mix with a product containing MCPA, may be applied in combination with other products registered for postemergence application in flax. Refer to the label of the tank mix partner for lists of other weeds controlled, rates of application, and use precautions.

Extreme growing conditions such as drought or near freezing temperatures prior to, at, and following time of application may reduce weed control and increase the risk of crop injury at all stages of growth. Only weeds that have emerged at the time of application will be affected. If foliage is wet at the time of application, control may be decreased.

## Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 60 days of harvest.
- Do not apply after flax has begun bolting as crop injury may occur if applied during the bloom period.
- Make one broadcast application per crop per year.
- Carefully follow rotational crop restrictions and other use precautions and limitations on the product label for Stinger HL.

### Conditions and Risks of Use for Special Local Need

**USE OF Stinger® HL HERBICIDE (THE “PRODUCT”) ON FLAX (THE “CROPS”) FOR THIS SPECIAL LOCAL NEED MAY RESULT IN CROP INJURY, CROP YIELD REDUCTION AND/OR CROP LOSS AS FURTHER DISCUSSED BELOW. READ AND UNDERSTAND THESE CONDITIONS AND RISKS OF USE FOR SPECIAL LOCAL NEED BEFORE USING THE PRODUCT ON THE CROP.**

This Product is available for use in the manner described in this Special Local Need Labeling on the basis that, in the sole opinion of the user, the benefits and utility derived from the use of the Product on the Crops outweigh the potential risk of Crop injury, Crop yield reduction or Crop loss. The decision to use this Product in the manner described in this Special Local Need Labeling must be made by each individual user on the basis of anticipated benefits versus (i) the potential risk of Crop injury, Crop yield reduction and Crop loss, (ii) the severity of the target pest infestation, (iii) the cost and availability of alternative pest controls and (iv) any other relevant factors.

**By purchasing the Product for use, or using the Product, in the manner described in this Special Local Need Labeling, you acknowledge and accept that to the extent permitted by law:**

- (1) you assume all risk of Crop injury, Crop yield reduction and Crop loss;
- (2) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any representations or recommendations regarding the use of this Product on the Crops other than the statements on this Special Local Need labeling;
- (3) Corteva Agriscience does not make, and does not authorize any agent or representative to make, any warranties, express or implied, with respect to the use of the Product on the Crops and disclaim all warranties, expressed or implied, including any implied warranty of merchantability;
- (4) Corteva Agriscience disclaims all liability for any damages, losses, expenses, claims or causes of actions arising out of or relating to Crop injury, Crop yield reduction and/or Crop loss;
- (5) these Conditions and Risks of Use for Special Local Need supersede any contrary representations or recommendations by Corteva Agriscience, or its respective agents or representatives, and any provisions in or on any Product literature or labeling including any provisions on the label affixed to the Product container.

**If these Conditions and Risks of Use for Special Local Need are not acceptable, the unopened Product may be returned to the seller for a refund or used for a different labeled use in accordance with the label affixed to the Product container.**

**These Conditions and Risks of Use for Special Local Need are required by Corteva Agriscience and not specified by the US EPA or the State of Montana.**

™® Trademarks of Corteva Agriscience and its affiliated companies

R469-007

Accepted: \_\_\_/\_\_\_/\_\_\_

Initial Printing