


2021

Cooperative Agricultural Pest Survey Report



Montana Department of Agriculture

Christy Clark, Director

Agricultural Sciences Division

Ian Foley, Administrator, SPRO

Commodity Services Bureau

Bureau Chief (Currently Vacant)

Layla Dunlap (2021)

State Survey Coordinator, SSC

Frank Etzler

Natural Resource Section

Alyssa Piccolomini, Carson Thomas

Agriculture Specialists

Ryan Solberg, Dawn Bales, Robyn Cassel, Taylor Cline, Jeff Drummond, Margaret Misner, Sean Mulla, Lori Vance

Produce Specialist

Larry Krum

Plant Pest Specialist

Pat Wherley (Ret.)

Agriculture Seasonals

Dan Poff, Therese Sutton

Agricultural Services Bureau

Bureau Chief

Beth Eiring

Noxious Weed Section

Jasmine Chaffee, Greta Dige, Andrew Lam, Josh Wagoner

Cooperators

USDA APHIS Plant Protection and Quarantine

USDA Forest Service

Montana State University Extension

Montana Department of Natural Resources and Conservation

US Department of the Interior

Montana Urban and Community Forestry Association

This report was compiled by Frank E. Etzler. Questions or comments can be addressed to the Montana Department of Agriculture Commodity Services Bureau at 302 North Roberts, Helena, Montana, 59601, or by phone at 406-444-9066, or email at agr@mt.gov

TABLE OF CONTENTS

Introduction to the CAPS Program _____	3
LDD Moth <i>Lymantria dispar</i> (L.) survey _____	4
Karnel Bunt (KB) <i>Tilletia indica</i> Mitra survey _____	6
Forest Pest Survey – Pest Detection _____	7
Small Grains Commodity Survey – Pest Detection _____	12
Exotic Woodborer and Bark Beetle Survey (EWBB) _____	14
Nursery Pest Survey _____	16
Pest Alert: Brown Marmorated Stink Bug (BMSB) _____	17
Terrestrial Mollusk Survey and Eastern Health Snail Update _____	18
National Honey Bee Survey _____	20
Japanese Beetle (JB) Surveys _____	22
National Agriculture Pest Information System (NAPIS) Survey Report	23

INTRODUCTION TO THE CAPS PROGRAM

The Cooperative Agricultural Pest Survey (CAPS) program conducts science-based national and state surveys targeted at specific exotic plant pests, diseases, and weeds identified as threats to U.S. agriculture and/or the environment. These activities are accomplished primarily under USDA funding that is provided through cooperative agreements with state departments of agriculture, universities, and other entities. Surveys conducted through the CAPS Program represent a second line of defense against the entry of harmful plant pests and weeds. These surveys enable the program to target high-risk hosts and commodities, gather data about pests specific to a commodity, and establish better baseline data about pests that were recently introduced in the United States. The mission of the CAPS program is to provide a survey profile of exotic plant pests in the United States deemed to be of regulatory significance through early detection and surveillance activities.

The Cooperative Agricultural Pest Survey is a nationwide survey effort initiated by the USDA Animal Plant Health Inspection Service (APHIS) Plant Protection and Quarantine (PPQ), to detect and/or monitor the spread of invasive plant pests. To achieve this goal, the USDA APHIS PPQ enlists the assistance of state cooperators. In Montana, state cooperators are coordinated through the Montana Department of Agriculture (MDA), and include not only the Department of Agriculture, but also Montana State University, the Montana Department of Natural Resources and Conservation, USDA Forest Service, and others.

CAPS Program Internet Resources

CAPS Website: <https://caps.ceris.purdue.edu/home>

National Agricultural Pest Information System (NAPIS): <http://pest.ceris.purdue.edu/>

Hungry Pests: <http://www.hungrypests.com/>

Montana Wood Boring Insect Project: <http://mtent.org/projects/woodboring/index.html>


LDD MOTH (LDDM) DETECTION SURVEY

Lymantria dispar (L.)

The European strain of the LDD moth (formerly Gypsy Moth) (*Lymantria dispar* (L.)) was initially introduced into the eastern United States in the mid-1800s. It established rapidly and became a serious defoliating forest pest. Over 500 susceptible host plants have been identified. Most are deciduous trees and shrubs, but older LDD moth larvae will also consume pine and spruce. In Montana, aspen and western larch are of particular importance as potential native tree host of the LDD moth, especially in the western half of the state. Most landscape plants, urban trees and shrubs throughout the state would also be subject to LDDM defoliation.

Females of the European strain are flightless but crawl actively as they seek out oviposition sites. The egg masses are brownish clumps covered with scales and hairs, and have been found on Christmas trees, boats, RVs, outdoor furniture, firewood, and virtually any other object that might be left outdoors in an infested area. They are thus readily transported to new areas by human activity. The LDD moth is the most destructive forest pest in the eastern United States and large areas of the northeastern and midwestern US are under a federal quarantine to prevent the spread of this pest. There are several other sub-species of closely related LDD moths from Asia that are not known to occur in North America but are attracted to the same pheromone lure. Asian LD Moth (ALDM) pest pressure has increased in recent

years due to increased populations in their native range and changes in international shipping logistics.



http://www.aphis.usda.gov/plant_health/plant_pest_info/gypsy_moth/downloads/gypmoth.pdf

There have been several detections of ALDM sub-species adults in the Pacific Northwest. In this sub-species, the female moths can fly and the caterpillars are more likely to feed on coniferous trees.

There have been several positive gypsy moth traps in Montana counties in recent years: Cascade (1989, 1990), Gallatin (1988), Glacier (2001, 2003, 2007, and 2008), Flathead (2019), Lewis and Clark (1988), Lincoln (2009), Liberty (1992), Missoula (1996), Park (2001), Yellowstone (1993 and 2011). Given the distance between Montana and the quarantined portions of the US and eastern Canada, it is almost certain these introductions were the result of human activity. Additional support for this is that most, if not all, of these counties are major recreational destinations for the entire U.S. Isolated detections result from the movement of egg masses and pupae on contaminated vehicles and equipment or adult moths “hitchhiking” with vehicles or other conveyances.



Male LDD Moth. Traps are baited with female sex-pheromone lures and only attract males.

In Montana, responsibility for the trapping of gypsy moths is a multi-agency cooperative effort between the USDA APHIS PPQ, the Montana Department of Agriculture (MDA), the Montana Department of Natural Resources & Conservation (DNRC), and the USDA Forest Service (USDA FS). In 2021, USDA APHIS PPQ placed traps in the eastern portion of the state, MDA placed traps in the western portion of the state, and DNRC placed traps in Missoula county. The USDA Forest Service, US Fish and Wildlife Service, Bureau of Indian Affairs, and other agencies coordinate trapping at a large number of campgrounds and other public recreation areas. The Department of the Interior placed traps in Glacier and Yellowstone National Parks. All traps were placed by early June, and checked throughout the summer.



LDD moth caterpillar. Via CT Dept. of Energy and Environmental Protection

RESULTS: 150 traps were placed by MDA, 50 traps by DNRC, 319 traps by USDA APHIS PPQ, and 306 by USFS in 2021. One trap in Fergus county, placed by USDA APHIS PPQ, was positive for the presence of LDD moth. A delimitation survey, conducted by USDA APHIS PPQ, is planned for the location in 2022. No other traps were positive.

Karnal Bunt Detection Survey

Tilletia indica Mitra


Karnal bunt (KB) is a fungal disease that affects wheat, durum wheat, and triticale. The disease was discovered near Karnal, India in 1931, hence the name. The first detection of KB in the United States was in Arizona in 1996, in durum wheat seed. Subsequently, the disease was found in portions of Southern California and Texas. The disease has never been detected in Montana field production. KB thrives in cool, moist temperatures as the wheat is starting to head out.

Karnal bunt spores are windborne and can spread through the soil. Spores have the ability to survive within the soil for several years. Grain can also become contaminated through equipment. Therefore, controlling the transportation of contaminated seed is essential in preventing the spread to Montana production areas. In addition, early detection is essential if any type of control or eradication is to be attempted. Montana's participation in the annual karnal bunt survey is part of the early detection grid set out across the United States.

RESULTS: Montana continued to sample for KB during the 2021 harvest. A total of 85 samples were collected from 16 counties across Montana. The APHIS Arizona State Plant Health Director's (SPHD) office Karnal bunt lab conducted the testing. All samples tested negative for the presence of KB. This sampling is critical for wheat growers in Montana. It confirms our wheat is free from KB, ensuring access to international export markets.



Bunted Wheat Credit: R. Duran, Washington State University www.forestryimages.org




Teliospores of *Tilletia indica* (Karnal bunt of wheat) showing surface ornamentation patterns. Credit:

EPPO.

Forest Pest Survey Pest Detection Survey

Forest land occupies an estimated 25.9 million acres in Montana (USDA 2019). Seventy-three percent (18.9 million acres) is publicly owned and under the jurisdiction of federal and state agencies (MT DNRC 2010, USDA 2019). Ecologists recognize 10 different major forest types in Montana. Douglas-fir (*Pseudotsuga menziesi* var. *glauca*), lodgepole pine (*Pinus contorta*), and ponderosa pine (*Pinus ponderosa*) predominate on the forest landscape and are the most commercially important species (MT DNRC 2010). Montana forests provide a wide variety of commercial and recreational benefits that are at risk from both native and invasive forest pests.



Pine Sawfly Detection Survey *Diprion pini* (L.)

Diprion pini is considered one of the most serious pests of pine in Russia, Ukraine, and Belarus. In Russia, outbreaks usually occur in 3–6 year intervals after hot and dry summers (Sharov 1993). Larvae are gregarious feeders and attack the shoots as well as mine the needles from the side. Larvae may also eat the bark of the shoots and may sometimes consume the shoots completely. Sawflies, including *D. pini*, highly prefer pine stands on infertile and well-drained soils as well as stands that are affected by unfavorable climatic or anthropogenic factors (Augustaitis 2007).



A native sawfly, *Neodiprion* sp. (male), caught in a pine sawfly trap. Photo: I. Foley

RESULTS: 25 pine sawfly traps were placed by MDA in 2021. 52 traps were placed by USDA PPQ in 2021. All traps were negative.

Masson Pine Moth (MPM) Detection Survey

Dendrolimus punctatus Walker, 1855

Dendrolimus punctatus, the Masson pine moth, occurs in China, Japan, Taiwan, and Vietnam. It is one of the most important forest pests in Southeast Asia, and defoliation of pine plantations during larval outbreaks causes significant economic loss. The primary host is *Pinus massoniana*, but larvae have been recorded feeding on a wide variety of other *Pinus*. Up to five overlapping generations per year are possible, with fewer generations in northern regions (Gilligan and Passoa 2014a). Because of this potential for serious damage to Montana's pines, the USDA monitors for this pest throughout the state.

RESULTS: 52 Masson pine moth traps were placed by USDA PPQ in 2021. All traps were negative.



Masson Pine Moth from Hong Kong, China. © 2019 Young Chan

Pine Beauty Moth (PBM) Detection Survey

Panolis flammea (Denis & Schiffermüller, 1775)

Panolis flammea is a colorful, nocturnal moth in the family Noctuidae native to Europe and Asia (Novak 1976). The species is considered a severe defoliator of *Pinus* species throughout many parts of Europe. Outbreaks in pine plantations in the United Kingdom and Continental Europe have caused damage to thousands of acres and resulted in significant mortality (Gilligan and Passoa 2014b). In the UK, adults are active from March through May. For Montana, lodgepole pines are especially at risk, as *P. flammea* has attacked these trees when planted in Scotland (Bradshaw et al. 1983, Sukovata et al. 2003). Monitoring for this species through CAPS pheromone traps and limiting the potential for establishment helps to ensure that Montana's characteristic pines are protected from possibly severe defoliation and any resulting impacts that could follow.

RESULTS: 25 traps were placed by the MDA and 52 traps were placed by the USDA PPQ in 2021. All traps were negative.



Panolis flammea resting, UK. © 2011 Malcom Storey

Pine Processionary Moth (PPM) Detection Survey

Thaumetopoea pityocampa (Denis & Schiffermüller, 1775)

Thaumetopoea pityocampa, the Pine Processionary Moth, is a moth in the family Notodontidae native to the Mediterranean (Southern Europe, North Africa, and the Middle East). Larvae in the genus are known to form long lines, or processions, on their way to new feeding locations. The larvae are the main defoliators, feeding upon pine needles, and are a major economic pest of coniferous forests in southern Europe (Bonnet et al. 2008). Additionally, the larvae (caterpillars) are covered in long urticating setae (hairs) that contain a toxin, thaumetopoein. These hairs can lead to severe skin dermatitis and allergic reactions in both people and animals upon contact with the larvae, the nests that the larvae build, or wind-blown loose hairs (Gilligan et al. 2014). Adults are active from May to September.



Thaumetopoea pityocampa in typical resting position.

Photo by Entomart.


RESULTS: 25 traps were placed by the MDA in 2021. All traps were negative.

Scots Pine Blister Rust (SPBR) Detection Survey

Cronartium flaccidum (Alb. & Schwein.) G. Winter 1880

Scots Pine blister rust (SPBR) is a heteroecious rust fungus native to Europe and Asia. The fungus can cause spotting on needles, and cankers on the stem that produce resin. These cankers can eventually lead to girdling of the stem, resulting in the death of the top or entire tree (Smith et al. 1988). Introduction of this disease to North America could have a major impact on Montana's forests and the health of our ecosystem.

RESULTS: 45 locations were visually surveyed for symptoms of the disease (Pine Detection and EWBB). No symptomatic plants were discovered. All surveys are part of the effort that goes into protecting Montana's forests for the future.




Pine Commodity Survey			
Target Species	Common Name	Approved Method	Sites
<i>Cronartium flaccidum</i>	Scots Pine Blister Rust	Visual	25
<i>Diprion pini</i>	Pine Sawfly	Delta trap/lure 28 days	77
<i>Dedrolimus punctatus</i>	Masson Pine Moth	Delta trap/lure 21 days	52
<i>Panolis flammea</i>	Pine Beauty Moth	Bucket trap/ lure 42 days	77
<i>Thaumetopoea pityocampa</i>	Pine Processionary Moth	Delta trap/lure 28 days	25

Small Grains Commodity Survey

Pest Detection Survey

Montana consistently ranks in the top 5 states for wheat and barley production in the United States. In 2020, Montana was ranked third of all US states for total wheat production and ranked second for Barley production. The 2021 growing season was severely affected by drought, with estimated yields for winter and durum wheat the lowest since 1988. Since wheat and barley are large portions of Montana’s Agricultural production, protecting these crops is essential.



In 2007, the USDA published guidelines for a small grains commodity based survey. The idea behind commodity based surveys is to target export commodities rather than individual pests. Here, multiple survey methods are used to take samples from a single commodity or group of similar commodities over a longer period of time. In the small grains survey, MDA used sweep net samples, visual surveys, soil samples for nematodes, and whole plant samples for diseases. This methodology allows the survey to maximize the potential for pest detection and minimize the cost compared to several different surveys for individual pests.

The 2020 small grains survey targets nine (9) different types of pests (see table below). These pests include 5 arthropods, 2 mollusks, and 2 nematodes. In addition to the

9 exotic pests, samples were also screened for the cereal leaf beetle and a number of other economically important nematodes and plant diseases.

RESULTS: During the 2021 survey, 23 sweep net samples were collected and 46 visual surveys were taken. Soil samples for nematode detection analysis were sent to the University of Nebraska in Lincoln. No suspect target pests were detected in any of the samples.

Whole plant samples were screened for disease by the Schutter Diagnostic Lab at Montana State University. The following non-regulated pathogens were detected: *Cochliobolus* Root Rot; *Fusarium* Root Rot; *Pyrenophora teres*, net blotch, spot form and net form; *Rhizoctonia* Root Rot; and *Zanthomonas* sp., bacterial wheat streak.

Small Grain Commodity Survey			
Target Species	Common Name	Approved Method	Sites
<i>Anguina tritici</i>	Wheat Seed Gall Nematode	Visual	23
<i>Ceruella virgata</i>	Maritime Garden Snail	Visual	23
<i>Cochlicella</i> spp.	Pointed Snail	Visual	23
<i>Diabrotica speciosa</i>	Curcubit beetle	Visual	23
<i>Eurygaster integriceps</i>	Sunn Pest Cereal Bug	Visual	23
<i>Helicoverpa armigera</i>	Old World Bollworm	Bucket Trap/ Lure 28 days	23
<i>Laodelphax striatellus</i>	Small Brown Planthopper	Yellow Sticky Card	23
<i>Meloidogyne artiellia</i>	Root-knot Nematode	Soil Sample	23
<i>Spodoptera littoralis</i>	Egyptian Cottonworm	Bucket Trap/ Lure 84 days	23



Small Brown Planthopper, *Laodelphax striatellus*.

© Paul Langolis, Museum Collections: Cicadas, Planthoppers, & Allies, USDA APHIS PPQ, Bugwood.org.

Exotic Woodborer and Bark Beetle (EWBB) Detection Survey – PPA 7721

Wood boring insects are some of the most dramatically destructive invasive species introduced into the forest and urban landscapes of the United States. These include notorious pests, such as the Asian Longhorned Beetle and the Emerald Ash Borer. Some native wood boring insects, such as the mountain pine beetle, also cause significant damage to Montana’s forest resources. However, the threat of exotic wood borers is significant for Montana’s agriculture, wood products, tourism, and recreation industries, as these exotic pests are freed from predators and diseases found in their native ranges.

The exotic woodborer and bark beetle (EWBB) survey targets primarily three groups of insects; longhorned beetles (Cerambycidae), bark beetles (Curculionidae: Scolytinae), and wood wasps (Siricidae). Within these groups, six species were specifically targeted in 2021, including Asian Longhorned Beetle and European Spruce Bark Beetle. This survey is conducted by using Lindgren funnels and panel traps baited with various ultra-high release (UHR) ethanols, bark beetles pheromone, and plant volatile lures. Funnel traps also have passive flight intercept capabilities, and the resulting trap catches include many native wood-boring beetles and a range of non-target families. While not specifically targeted, flight intercepts do capture beetles in the family Buprestidae and have the potential to trap exotic buprestids such as the Emerald Ash Borer.

In 2021, 40 funnel traps and 20 vane traps were placed and monitored across the state cooperatively by MDA and Montana State University. Trap sites focused on forested areas near the Canadian border, recreation sites with campgrounds, and high traffic tourism areas.

Exotic Woodboring Beetle Survey			
Target Species	Common Name	Approved Method	Sites
<i>Anoplophora glabripennis</i>	Asian Longhorned Beetle	Visual	20
<i>Cronartium flaccidum</i>	Scots Pine Blister Rust	Visual	20
<i>Hylobius abietis</i>	Large Pine Weevil	Multifunnel Trap/ Lure EtOH, a-pinene UHR, Monochamol	20
<i>Ips sexdentatus</i>	Sixtoothed Bark Beetle	Multifunnel Trap/ Lure Ips, 3 dispenser	20
<i>Ips typographus</i>	European Spruce Bark Beetle	Multifunnel Trap/ Lure Ips, 3 dispenser	20
<i>Monochamus urussovii</i>	Black Fir Sawyer	Multifunnel Trap/ Lure EtOH, a-pinene UHR, Monochamol	20
<i>Trichoferus campestris</i>	Velvet Longhorned Beetle	Cross Vane Trap/ <i>Trichoferus campestris</i> Lure	20

RESULTS: No target species (see table above) were collected. Due to the extreme drought this year in Montana and associated wildfires, a few localities were unable to be reached due to fires. These localities are planned to be surveyed in the future to help track changes to beetle fauna after burns and continue monitoring for potential invasive species.

Cerambycidae: There are 152 species of longhorned beetles recorded from Montana (Hart et al. 2013). The most common species in funnel traps include *Acmaeops proteus* (Kirby), *Arhopalus asperatus* (LeConte), *Asemum striatum* (Linnaeus), *Monochamus scutellatus* (Say), *Neandra brunnea* (Fabricius), *Neospondylis upiformis* (Mannerheim), *Rhagium inquisitor* (Linnaeus), *Tetropium velutinum* LeConte, and *Xylotrechus longitarsis* Casey.

Scolytinae: There are approximately 100 species of bark beetles recorded from Montana (Gast et al. 1989, NAPIS 2012).

Montana Wood Boring Insect Project

Montana State University through the Montana Agricultural Experiment Station (MAES) and Montana Entomology Collection (MTEC) has developed an online portal for the “Montana Wood Boring Insect Project”. This website contains county level distribution data and images of all of the long-horned and metallic wood boring beetles known to occur in Montana. Many of the non-target species collected through the CAPS program have been incorporated into this project and are maintained in the MTEC. The project website can be found at: <http://mtent.org/projects/woodboring/index.html>



Entomology Group / Montana Entomology Collection (MTEC) / Wood Boring Insects

Wood Boring Insects of Montana



Wood Boring Insect Families

- [Horntails \(Siricidae\)](#)
- [Longhorn Beetles \(Cerambycidae\)](#)
- [Metallic Flathead Borers \(Buprestidae\)](#)
- [Powder Post Beetles \(Bostrichidae\)](#)
- [Bark Beetles \(Scolytinae\)](#)

A Montana Agriculture/ Experiment Station project in cooperation with [USDA, APHIS, Plant Protection and Quarantine](#), [Montana Department of Agriculture](#), [Dept. of Natural Resources and Conservation, Forestry Division](#), and [Montana 4-H](#)

Nursery Pest Detection Survey – PPA 7721

The nursery industry is important as it allows people to cultivate gardens for beauty and to produce food for personal consumption. However, due to the plant trade between states, this could create artificial pathways for the introduction of potential pests. By monitoring these pathways, we can ensure that Montana’s green industry is free from any regulated pests and is protected from accidental introductions. In Montana, over 1000 businesses hold nursery licenses, so ensuring that they are protected is crucial to our green industry.

Part of the survey is to also survey the western portion of the state for Asian Giant Hornet (AGH, *Vespa mandarinia* Smith, see cover). This species was first discovered in Washington State in 2019. Since then, eradication efforts have been conducted to limit the spread and to attempt to eliminate the species from North America. The MDA’s survey work ensures that AGH doesn’t spread artificially or naturally into Montana, which could have serious impacts on Montana’s Apiaries.

Another pest surveyed for is the Tomato Leaf Miner (*Tuta absoluta* Meryrick). This moth is native to South America and has spread globally in recent years, and can oviposit on any plant in Solanaceae. Within Montana, the crop of concern are potatoes, which are grown for seed in Montana. Serious impacts could occur to Montana’s Seed potato industry if this species was to become established.

Other pests, such as the Spotted Lanternfly, African Giant Land snail, and the Christmasberry webworm, were monitored for as well, but economic impacts of these species is expected to be limited in Montana.

RESULTS: 25 nursery locations were sampled and inspected in Montana. No regulated pests were discovered.



Tomato leafminer, *Tuta absoluta*. Credit: Marja van der Straten, NVWA Plant Protection Service, Bugwood.org.

PEST ALERT: Brown Marmorated Stinkbug (BMSB)

Halyomorpha halys Stål



Brown Marmorated Stinkbug. Credit: Steven Valley, Oregon Department of Agriculture, Bugwood.Org

The Brown Marmorated Stinkbug (BMSB, *Halyomorpha halys* Stål) was first discovered in the US in Pennsylvania in 1998. Since then, the species has spread across North America. In January 2021, BMSB was discovered at a local residence in Billings, Montana. In May 2021, a second specimen was found flying within a personal vehicle in Flathead County. These two localities suggest a much larger distribution than currently known.

BMSB is a pest of concern as it has a large host range, and could affect crops, such as corn, and specialty crops, such as Flathead cherries, and personal gardens. BMSB overwinters in large aggregates, and is often considered a nuisance pest of residential homes in areas where it is established.

The MDA, along with Montana State University Extension services, have planned surveys across Montana in 2022 to determine distribution of BMSB within the state. Preliminary studies conducted by a professor out of Minnesota, who is an expert in BMSB, have confirmed that there is an established population in Billings, Montana.

Terrestrial Mollusk Survey and Eastern Health Snail (EHS) Update

PPA 7721

BACKGROUND

Snail samples collected in Cascade County in late July of 2012 were confirmed as Eastern Health Snail (EHS), *Xerolenta obvia*, one of twelve USDA listed invasive terrestrial snails of national concern. The Montana Department of Agriculture and Montana PPQ conducted survey work in August and September of 2012 to delimit the infested area, determine whether eastern health snail was present in grain and alfalfa production areas in the state, and to support export of Montana agricultural commodities and products. Survey work confirmed the presence of snails in the Belt area along transportation corridors, residential areas, rangeland, hay fields, and yards. Extensive survey work outside the infested area showed snails were not yet present in grain production areas. Through discussion with individual Belt area landowners and residents, it was determined the snails have been present in the area for at least 25 years, perhaps much longer. Pathways of introduction include rail, mining, travel, and trade/commerce. There is a strong correlation between rights-of-way activities and local distribution of the snail. In 2013, two additional populations of *Xerolenta obvia* were confirmed in Cascade County (in the city of Great Falls and near Monarch).

MITIGATION EFFORTS

Since 2018, laboratory studies on Eastern Health Snail have been conducted at Montana State University by Jennie Birdsall, Jeff Littlefield, and Annie deMeij. These studies have focused on the development of EHS and determining suitable hosts for feeding. Studies are also being conducted to determine possible biological control of EHS.

SURVEY

Since initial detection, surveys for invasive terrestrial mollusks have occurred every year. Survey sites included high-risk transportation areas, recreational areas, and nurseries. Survey work was focused on presence or absence of snails and no attempt was made to quantify the snail population. Survey work appears to indicate that snails have not spread beyond the infested boundaries identified in 2012. It remains important to conduct survey work in the future to monitor the snail population in the Belt area and determine presence or absence in other areas to support Montana's export markets.



Cochlicella sp. on grain



Maritime garden snail, *Cerņuella virgata*

Mollusks have only recently been identified as a threat in Montana. Movement of various materials protected by solid wood packing material into and through Montana increases the risk of introduction of pests – not only through standard commerce, but also through the movement of materials from the seaport inland. Interstate 90, a major route across the U.S., travels the entire width of Montana, from a point just west of Missoula to east of Glendive. The Montana “banana belt,” a region of milder climate, runs from the Flathead Valley to the Bitterroot. This area has experienced a rapid influx of people and an increase in the building of higher-value homes, with rates exceptionally high in 2020 and 2021. These properties often include imported materials such as tile, marble, and wood.

The entire state of Montana is a mecca for recreation including activities of all types. All of these serve as routes of entry into the state for organisms such as the various Veroncelid snails, as well as *Monacha* spp., *Cerņuella* spp., and *Cochlicella* spp. These snails could, if established, not only out-compete native species, but also eliminate portions of the food web that are currently supporting the state’s famous trout fisheries, become mechanical obstacles to field crop harvest, and directly damage desirable plant species including wheat.

RESULTS: No additional invasive mollusk species or EHS localities were discovered in 2021.



Eastern Heath Snail, *Xerolenta obvia*. Photo by Ian Foley

2021 National Honey Bee Survey

In an average year, Montana has about 275,000 beehives, of which about half are migratory. Montana has about 550 registered beekeepers, about one-fifth of whom are commercial operators. Most of these provide commercial pollination services outside Montana. Migratory beekeepers typically travel to California in the early spring for almond pollination, then move to fruit crops in Washington and Oregon before moving back to Montana for the summer. Ranked by revenue, beekeeping is the 10th largest agricultural industry in Montana; pollination fees make up the majority of that income.

Pests of honey bees are a serious threat to the agricultural economy of Montana and to the states where Montana-based bees provide pollination. USDA estimates honey bee pollination adds approximately \$15 billion to the value of American agriculture. In 2006 beekeepers began reporting unexplained and unexpected losses of 30% or more of their hives. What eventually came to be called “colony collapse disorder” (CCD) was characterized by the rapid disappearance of worker bees from apparently healthy hives. Despite a considerable increase in honey bee research, the cause of colony collapse remains unknown, and unexplained losses continue at about 30% per year. Recent research has focused on pathogen load, pesticide exposure, stress, and habitat modification. The current hypothesis for CCD is that it may be a symptom of a complex of factors.

In 2009 the USDA-APHIS initiated the National Honey Bee Pests and Diseases Survey in all 50 states. The primary objectives of the survey are to confirm the absence of tropical bee mites in the genus *Tropilaelaps*, the absence of the Asian honey bee *Apis ceranae*, and the absence of slow paralysis virus (a honey bee disease associated with *A. ceranae*). Secondary objectives include evaluating the overall health of the apiaries sampled to establish a baseline for future research. Samples submitted from the survey will be evaluated for their mite loads (*Varroa*, tracheal mites, and other parasitic mites) and the degree to which viruses and other pathogens are present (particularly *Nosema ceranae*, a more virulent *Nosema* species associated with tropical honey bees). Viruses are identified at the molecular level by the USDA “bee lab” in Beltsville, MD.



Montana bee yard. Photo: A. Piccolomini



Varroa mites on a drone pupa. Photo I. Foley

RESULTS: 25 National Honey Bee Survey (NHBS) samples were collected in 2021 and submitted to laboratories at the University of Maryland. The 2021 season is the first season where all 25 samples were taken in the same season, thanks to combined efforts of the State Entomologist and Apiary Tech. Due to the ongoing effects of the COVID-19 pandemic, some results are still pending. Nosema Disease (*Nosema* spp.), Chronic Bee Paralysis Virus, Deformed Wing Virus, Israeli Acute Paralysis Virus, Lake Sinai Virus-2, and Varroa Destructor Virus were all detected in at least one sample. European Foulbrood was discovered in one of the NHBS samples. American Foulbrood and Small Hive Beetle were discovered at a single location and the location was quarantined, with eventual voluntary destruction of infected material.



Honey bee Workers and Queen. Photo A. Piccolomini

Japanese Beetle (JB) Surveys

Popillia japonica Newman



Japanese Beetle, *Popillia japonica* Newman, was first discovered in North America in 1916. Since then, it has spread throughout much of eastern North America. Japanese Beetle (JB) was first discovered in Billings, Montana in 2001. In 2013, nursery stock infested with JB was brought into Montana, affecting 17 nurseries across the state. Due to rapid action by the nurseries and MDA, these localities were negative in 2014. The only area of Montana with a consistent population of JB is Billings, Montana, although the population has moved within city limits since first detection.

2021 also saw the implementation of a Specialty Crop Block Grant for community outreach and pesticide distribution for JB in Billings. This outreach effort aims to educate the public about JB and to provide tools for them to begin public control efforts to reduce the local JB population.

RESULTS: The MDA put out over 130 traps, focusing on Flathead, Sanders, Ravalli, and Yellowstone Counties. The USDA placed 32 traps at airports in Cascade, Flathead, Gallatin, Lewis and Clark, Missoula, and Silver Bow Counties to ensure no accidental introductions to the state via air travel. In Billings (Yellowstone Co.), three trap locations were positive for JB, with a total of 1971 beetles collected from July to September across all three traps.

National Agricultural Pest Information System (NAPIS)

2021 Summary Report

Pest Common	Pest Scientific	Data Source	Counties	Positives	Negatives	Total
Acute Bee Paralysis (ABPV)	<i>Aparavirus Acute Bee Paralysis</i>	State Ag Dept.	10	0	13	13
American Foulbrood	<i>Paenibacillus larvae larvae</i>	State Ag Dept.	10	1	13	14
Asian Giant Hornet	<i>Vespa mandarinia</i>	State Ag Dept.	11	0	23	23
Asian Honeybee	<i>Apis ceranae</i>	State Ag Dept.	10	0	13	13
Asian Longhorned Beetle	<i>Anoplophora glabripennis</i>	University/Extension	8	0	20	20
Black Fir Sawyer	<i>Monochamus urussovii</i>	University/Extension	8	0	20	20
Black Fir Sawyer	<i>Monochamus urussovii</i>	USDA APHIS	14	0	53	53
Chalkbrood	<i>Ascosphaera apis</i>	State Ag Dept.	13	0	17	17
Christmasberry Webworm	<i>Cryptoblabes gnidiella</i>	State Ag Dept.	12	0	25	25
Chronic Bee Paralysis (CBPV)	Unassigned Chronic Bee Paralysis Virus	State Ag Dept.	13	0	17	17
Curcubit beetle	<i>Diabrotica speciosa</i>	State Ag Dept.	7	0	21	21
Deformed Wing Virus (DWW)	<i>Iflavirus Deformed Wing Virus</i>	State Ag Dept.	10	7	6	13
Eastern Heath Snail	<i>Xerolenta obvia</i>	State Ag Dept.	9	1	47	48
Eqyptian Cottonworm	<i>Spodoptera littoralis</i>	State Ag Dept.	8	0	23	23
European Foulbrood	<i>Melissococcus plutonius</i>	State Ag Dept.	10	1	12	13
European Spruce Bark Beetle	<i>Ips typographus</i>	University/Extension	8	0	20	20
Giant African Snail	<i>Lissachatina fulica</i>	State Ag Dept.	12	0	25	25
Hygromiid Snails	<i>Ceruellea spp.</i>	State Ag Dept.	9	0	48	48
Hygromiid Snails	<i>Monacha spp.</i>	State Ag Dept.	9	0	48	48
Israeli Acute Bee Paralysis (IAPV)	<i>Aparavirus Israeli Acute Paralysis</i>	State Ag Dept.	10	5	8	13
Japanese Beetle	<i>Popillia japonica</i>	State Ag Dept.	4	3	129	132
Japanese Beetle	<i>Popillia japonica</i>	USDA APHIS	6	0	32	32
Japanese Pine Sawyer	<i>Monochamus alternatus</i>	USDA APHIS	14	0	53	53
Karnel Bunt	<i>Tilletia indica</i>	State Ag Dept.	16	0	85	85
Kashmir Bee Virus (KBV)	<i>Aparavirus Kashmir Bee Virus</i>	State Ag Dept.	10	0	13	13
Lake Sinai-2	<i>Sinavirus Lake Sinai Virus-2</i>	State Ag Dept.	10	5	8	13
Large Pine Weevil	<i>Hylobius abietis</i>	University/Extension	8	0	20	20
Large Pine Weevil	<i>Hylobius abietis</i>	USDA APHIS	14	0	53	53
LDD Moth	<i>Lymantria dispar dispar</i>	State Ag Dept.	12	0	150	150
LDD Moth	<i>Lymantria dispar dispar</i>	State DNRC	1	0	50	50
LDD Moth	<i>Lymantria dispar dispar</i>	USDA APHIS	14	1	317	318
LDD Moth	<i>Lymantria dispar dispar</i>	USFS	30	0	306	306
Maritime Garden Snail	<i>Ceruellea virgata</i>	State Ag Dept.	9	0	71	71
Masson Pine Moth	<i>Dendrolimus punctatus</i>	USDA APHIS	14	0	52	52
Moku Virus (MKV)	<i>Iflavirus Mokus Virus</i>	State Ag Dept.	10	0	13	13
Nosema spores	<i>Nosema ceranae</i>	State Ag Dept.	10	8	5	13
Old World Bollworm	<i>Helicoverpa armigera</i>	State Ag Dept.	8	0	23	23
Parasitic Mite	<i>Trpilaelaps spp.</i>	State Ag Dept.	10	0	13	13

Pest Common	Pest Scientific	Data Source	Counties	Positives	Negatives	Total
Parasitic Mite Syndrome	Parasitic Mite Syndrome	State Ag Dept.	10	0	13	13
Pine Beauty Moth	<i>Panolis flammea</i>	State Ag Dept.	6	0	25	25
Pine Beauty Moth	<i>Panolis flammea</i>	USDA APHIS	14	0	52	52
Pine Processionary Moth	<i>Thaumetopoea pityocampa</i>	State Ag Dept.	6	0	25	25
Pine Sawfly	<i>Diprion pini</i>	State Ag Dept.	6	0	25	25
Pine Sawfly	<i>Diprion pini</i>	USDA APHIS	14	0	52	52
Pointed Snail	<i>Cochlicella spp.</i>	State Ag Dept.	9	0	71	71
Root-knot Nematode	<i>Meloidogyne artiellia</i>	State Ag Dept.	8	0	23	23
Sacbrood	<i>Morator aetatulas virus</i>	State Ag Dept.	10	0	13	13
Scots Pine Blister Rust	<i>Cronartium flaccidum</i>	State Ag Dept.	6	0	25	25
Scots Pine Blister Rust	<i>Cronartium flaccidum</i>	University/Extension	8	0	20	20
Sixtoothed Bark Beetle	<i>Ips sexdentatus</i>	University/Extension	8	0	20	20
Slow Bee Paralysis (SBPV)	<i>Iflavirus Slow Bee Paralysis</i>	State Ag Dept.	10	0	13	13
Small Brown Planthopper	<i>Laodelphax stiatellus</i>	State Ag Dept.	8	0	23	23
Small Hive Beetle	<i>Aethina tumida</i>	State Ag Dept.	10	1	13	14
Spotted Lanternfly	<i>Lycorma delicatula</i>	State Ag Dept.	12	0	25	25
Sunn Pest Cereal Bug	<i>Eurygaster integriceps</i>	State Ag Dept.	7	0	21	21
Tomato Leaf Miner	<i>Tuta absoluta</i>	State Ag Dept.	12	0	25	25
Varroa Destructor Virus (VDV)	<i>Iflavirus Varroa Destructor Virus</i>	State Ag Dept.	10	8	5	13
Velvet Longhorned Beetle	<i>Trichoferus campestris</i>	University/Extension	8	0	20	20
Wheat Seed Gall Nematode	<i>Anguina tritici</i>	State Ag Dept.	9	0	22	22
REPORT TOTAL				41	2333	2374

References Cited

- Augustaitis, A. 2007.** Pine sawfly (*Diprion pini* L.)- related changes in Scots pine crown defoliation and possibilities of recovery. Polish Journal of Environmental Studies 16: 363–369.
- Bonnet, C., J. C. Martin, and R. Mazet. 2008.** La processionnaire du pin. Stantari - Histoire naturelle et culturelle de la Corse , Kyrnos publications 14: 29–33. [In French]
- Bradshaw, J. W. S., R. Baker, C. Longhurst, J. C. Edwards and J. C. Lisk. 1983.** Optimization of a monitoring system for the pine beauty moth, *Panolis flammea* (Denis & Schiffermüller), using sex attractants. Crop Protection 2: 63–73.
- Gilligan, T. M. and S. C. Passoa. 2014a.** Screening aid: Pine lappets, *Dendrolimus* spp. Identification Technology Program (ITP), USDA-APHIS-PPQ-S&T, Fort Collins, CO. 7 pp.
- Gilligan, T. M. and S. C. Passoa. 2014b.** Screening aid: Pine beauty, *Panolis flammea* (Denis & Schiffermüller). Identification Technology Program (ITP), USDA-APHIS-PPQ-S&T, Fort Collins, CO. 5 pp.
- Gilligan, T. M., S. C. Passoa and F. Groenen. 2014.** Screening aid: Processionary moths, *Thaumetopoea* spp. Identification Technology Program (ITP), USDA-APHIS-PPQ-S&T, Fort Collins, CO. 6 pp.
- Montana Department of Natural Resources and Conservation (DNRC). 2010.** Montana’s State Assessment of Forest Resources: Base Findings and GIS Methodology. State of Montana. 27 pp.
- Novak, V. 1976.** Atlas of Insects Harmful to Forest Trees. Volume 1. Elsevier Scientific Publishing Company. Amsterdam, The Netherlands. 125 pp.
- Sharov, A. A. 1993.** Biology and population dynamics of the common pine sawfly, *Diprion pini* L., in Russia. pp. 409–430. *In*: M. Wagner and K. F. Raffa, eds. Sawfly Life History Adaptations to Woody Plants. Academic Press, Inc. San Diego, California.
- Sukovata, L., A. Kolk, J. Jaroszynska, U. Krajewska, A. Purzynska and V. Isidorov. 2003.** Host-tree preferences of the pine moth (Lepidoptera: Lasiocampidae) and pine beauty moth (Lepidoptera: Noctuidae) larvae in relation to needle quality. pp. 98–106. *In*: M. L. McManus and A. M. Liebhold, eds. Proceedings: Ecology, Survey and Management of Forest Insects; 2002 September 1-5; Krakow, Poland. Gen. Tech. Rep. NE-311. Newtown Square, PA: U.S. Dept. of Agriculture, Forest Service, Northeastern Research Station. .
- [USDA] Witt, C., J. D. Shaw, J. Menlove, S. A. Goeking, R. J. DeRose, K. A. Pelz, T. A. Morgan, S. W. Hayes. 2019.** Montana’s forest resources, 2006–2015. Resource Bulletin RMRS-RB-30. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 102 pp.